

09-21-00

526 Rec'd PCT/PTO 20 SEP 2000 PCT

FORM PTO-1390
(REV 5-93)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

GEI-078

U.S. APPLICATION NO (if known, see 37 CFR 1.5)

09/646763

INTERNATIONAL APPLICATION NO
PCE/FR99/00680INTERNATIONAL FILING DATE
3/23/99PRIORITY DATE CLAIMED
3/23/98

TITLE OF INVENTION

PHARMACEUTICAL HORMONAL COMPOSITION WITH SYSTEMIC ACTION

APPLICANT(S) FOR DO/EO/US

LANQUETIN et al

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
 2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
 3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
 4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
 5. ☐ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
 6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
 7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
 8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
 9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
 10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).
- Items 11. to 16. below concern other document(s) or information included:
11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
 12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
 13. ☒ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
 14. ☐ A substitute specification.
 15. ☐ A change of power of attorney and/or address letter.
 16. ☒ Other items or information: First page of PCT application, form 2038 (02-2000)

09/646763

430 Rec'd PCT/PTO 20 SEP 2000

US APPLICATION NO. 09/646763	INTERNATIONAL APPLICATION NO. PCT/FR99/00680	ATTORNEY'S DOCKET NUMBER GEI-078
-------------------------------------	---	---

17. <input checked="" type="checkbox"/> The following fees are submitted:				CALCULATIONS		PTO USE ONLY
Basic National Fee (37 CFR 1.492(a)(1)-(5)): Search Report has been prepared by the EPO or JPO..... \$830.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) \$640.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)).. \$710.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37CFR 1.445(a)(2)) paid to USPTO..... \$950.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)..... \$90.00				\$970.00		
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$ 970.00		
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$		
Claims	Number Filed	Number Extra	Rate			
Total Claims	18 -20 -	0	X \$22.00	\$ 0		
Independent Claims	1 -3 -	0	X \$74.00	\$ 0		
Multiple dependent claims(s) (if applicable)			0 + \$230.00	\$ 0		
TOTAL OF ABOVE CALCULATIONS =				\$ 970.00		
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28).				\$		
SUBTOTAL =				\$ 970.00		
Processing fee of \$130.00 for furnishing the English translation later the <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+ \$		
TOTAL NATIONAL FEE =				\$ 970.00		
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$		
TOTAL FEES ENCLOSED =				\$ 970.00		
				Amount to be:		
				refunded \$		
				charged \$		

- a. ☐ A check in the amount of \$_____ to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. _____ in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 02-2275. A duplicate copy of this sheet is enclosed.
 x Fee authorization form 2038(02-2000) enclosed herewith.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:
 Bierman, Muserlian and Lucas
 600 Third Avenue
 New York, N.Y. 10016



20311

PATENT TRADEMARK OFFICE

SIGNATURE

Charles A. Muserlian

NAME

19,683

REGISTRATION NUMBER

09/646763

430 Rec'd PCT/PTO 20 SEP 2000

GEI-078

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: :
Michel Lanquetin, et al. : Group:
Filed: Concurrently Herewith :
For: TOPICAL HORMONAL : Examiner :
COMPOSITION WITH SYSTEMIC :
EFFECT :

600 Third Avenue
New York, NY 10016
September 20, 2000

PRELIMINARY AMENDMENT

Asst. Commissioner For Patents
Washington, D.C. 20231

SIR:

Please amend this application as follows:

IN THE SPECIFICATION:

Page 1, before line 1, insert

--This application is a 371 of PCT FR99/00680 filed March
23, 1999.--

IN THE CLAIMS:

Claim 3, line 1, cancel "or claim 2"

Claim 4, line 1, cancel "any one of the claims 1 to 3" and

insert --claim 1--.

Claim 6, line 1, cancel "any one of the claims 1 to 5" and

insert --claim 1--.

Claims 7 and 8, line 1, cancel "any one of the claims 1 to 6"

and insert --claim 1--.

Claim 9, line 1, cancel "any one of the claims 1 to 8" and

insert --claim 1--.

Claim 11, lines 1 and 2, cancel "any one of the claims 1 to

10" and insert --claim 1--.

Claim 14, lines 1 and 2, cancel "any one of the claims 1 to

13" and insert --claim 1--.

Claim 17, line 1, cancel "any one of the claims 1 to 16" and

insert --claim 1--.

Please add the following claim:

--18. A method of systemically effecting correction of progesterone deficiency in premenopausal women comprising administering to premenopausal women in need thereof a systemically correcting amount of a composition of claim 1 to treat progesterone deficiency.--

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

Respectfully submitted,
Bierman, Muserlian and Lucas

By:

Charles A. Muserlian
Charles A. Muserlian #19,683
Attorney for Applicants
Tel.# (212) 661-8000

CAM:mlp

Encl.: Return receipt postcard

Variable	Mean		SD		t		p	
	Control	Case	Control	Case	Control	Case	Control	Case
Age	30.5	30.5	1.2	1.2	0.0	0.0	1.000	1.000
Gender	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Education	12.0	12.0	1.0	1.0	0.0	0.0	1.000	1.000
Marital status	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Occupation	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Religion	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Income	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Health status	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Family size	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Urban/rural	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Marital satisfaction	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental satisfaction	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal satisfaction	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental involvement	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal involvement	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental communication	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal communication	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental support	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal support	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental discipline	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal discipline	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental monitoring	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal monitoring	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental control	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal control	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental warmth	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal warmth	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental rejection	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal rejection	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental criticism	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal criticism	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental praise	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal praise	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental affection	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal affection	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental hostility	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal hostility	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental aggression	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal aggression	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental withdrawal	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal withdrawal	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Parental rejection	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000
Spousal rejection	1.0	1.0	0.0	0.0	0.0	0.0	1.000	1.000

[illegible]

6/pst

09/646763

TOPICAL HORMONAL COMPOSITION WITH SYSTEMIC ACTION

430 Rec'd PCT/PTO 20 SEP 2000

The present invention relates to the area of therapeutic chemistry and especially to the development of new galenic forms for application on the skin.

5

The present invention relates more particularly to galenic preparations whose active principle is a synthetic progestogen, intended to be applied to the skin in order to achieve a systemic hormonal effect in women before and after the menopause.

10 Thus, the invention relates to a topical hormonal composition with systemic action.

In particular, French Patent 2.271.833 describes hormonal compositions for the correction of progestogen deficiencies in premenopausal or menopausal women, intended for oral administration.

5

However, the oral route is not without certain drawbacks for natural progesterone, as well as for synthetic progestogens. On the one hand, it requires the administration of quite large doses in order to make up for degradation of the active principle during passage through the intestine and in the liver (called the "first passage" effect). On the other hand, it does not give constant plasma levels over time since oral administration is followed by a plasma peak during which the blood concentrations are raised temporarily.

20

Natural progesterone is sometimes administered percutaneously. This route only produces local effects, and does not permit remote impregnation of the target tissues, notably the uterine mucosa. This is because there is rapid degradation of the hormone by enzymes in the subcutaneous tissue, making it impossible to reach sufficiently high plasma levels to produce a systemic hormonal action.

25

Many synthetic progestogens have the same drawback and cannot be used percutaneously to achieve a systemic effect. The only exception is norethisterone acetate administered in patches.

30

The skin's function as a protective barrier against external aggressive agents makes it rather impermeable to numerous substances and only allows medicinal molecules to penetrate under

certain conditions: size and nature of the molecule, solubility, stability, nature of the vehicle containing the molecule, etc.

Thus, the release of an active principle from a vehicle and its penetration through the skin and as far as the blood or lymphatic circulation depend on numerous physicochemical and/or physiological parameters.

In the present invention, the very nature of the active principle (synthetic progestogen) presents the main obstacle to penetration through the skin: the main problem that arises is poor diffusion through the epidermis on account of its lipophilic character. The choice of vehicle used in the compositions will therefore have a considerable influence on percutaneous penetration and on the therapeutic activity of the molecule.

Thus, the topical compositions according to the invention permit a systemic effect by optimization of percutaneous passage of a synthetic progestogen derived from 19-nor progesterone.

The topical compositions according to the invention contain, as active principle, a synthetic progestogen derived from 19-nor progesterone and excipients that ensure optimum passage of the active principle through the skin.

The present invention relates more specifically to a topical hormonal composition with systemic effect for correcting progesterone deficiencies in premenopausal women and for hormone replacement in menopausal women, characterized in that it comprises:

- as active principle, a progestogen derived from 19-nor progesterone,
- a vehicle permitting systemic passage of the said active principle chosen from the group that includes a solubilizing agent, an absorption promoter, a film-forming agent, a gelling agent or their mixtures,

combined with or mixed with suitable excipients for production of a pharmaceutical form as a gel and/or a film.

The compositions according to the invention can therefore be in the form of a gel, a film-forming gel or a film-forming solution.

The progestogen derived from 19-nor progesterone used in the present invention is nomegestrol and/or one of its esters or ethers.

An example of nomegestrol ether is tetrahydropyranic ether of nomegestrol.

An example of nomegestrol ester is nomegestrol acetate, which is a synthetic progestogen that is active in oral administration, whose action comprises correction of gynaecological disorders caused by deficiency of luteinizing hormones.

5

Administered by means of compositions according to the invention, nomegestrol acetate is able to pass through the skin and enter the blood circulation to give plasma levels that can be detected by the methods used for assaying in biological media. The plasma concentrations observed are maintained at a plateau after cutaneous application because of the reservoir effect of the skin.

10

The plasma levels of nomegestrol acetate obtained with the compositions according to the invention are able to create a hormonal effect on tissues located far from the site of application, and especially on the endometrium.

15

Repeated administration of nomegestrol acetate produces a therapeutic action when it is given to premenopausal women suffering from symptoms connected with progesterone deficiency or to menopausal women undergoing oestrogen replacement therapy.

20

According to the invention, nomegestrol or one of its esters or ethers is present in a quantity varying from 0.05 to 1 wt.% of the total composition. Preferably, nomegestrol or one of its esters or ethers is present in an amount varying from 0.1 to 0.8 wt.% of the total composition. The topical compositions with systemic effect that are preferred according to the invention are those containing a quantity of nomegestrol or of one of its esters or ethers of 0.4 wt.% of the total composition.

25

The solubilizing agents and the absorption promoters have different modes of action but they both favour penetration of the active principle through the skin.

The solubilizing agents improve the solubility of the active principle and alter its affinity for the skin by acting upon the thermodynamic activity of the active molecule.

30

The absorption promoters lower the resistance to diffusion by modifying the structure of the cutaneous barrier.

However, there is no direct relation between improvement of solubility of the active principle in the vehicle and increased passage through the skin. In fact, the use of agents that improve the solubility of the active principle also increases its affinity for the vehicle and therefore generally lowers its diffusion through the skin.

- 5 Thus, for the solubility of the active principle in a vehicle to be total, there must be a certain affinity for the latter; however, it must not be too great, so that division of the active principle is oriented towards its diffusion through the skin.

According to the present invention, examples of suitable solubilizing agents are water, alcohols,
10 propyleneglycol, polyethylene glycol, polyethylene 20 sorbitan mono-oleate (marketed for example under the name Polysorbate 80 DF), a C₈/C₁₀ polyoxyethylene glycosyl glyceride (marketed for example under the trade-name Labrasol®) or their mixtures.

5 The solubilizing agent used is generally a mixture of solvents or of the aforementioned solubilizing agents, which, by synergistic action, is more effective than one of them used alone.

The solubilizing agent is preferably chosen from the group comprising water, alcohols, propyleneglycol, a C₈/C₁₀ polyoxyethylene glycosyl glyceride or their mixtures.

Thus, it will be possible to use, as solubilizing agent, a binary mixture of 95° ethanol and water, in which the percentage of 95° ethanol varies from 30 to 50%, and especially a binary mixture of
20 95° ethanol and water in which the percentage of 95° ethanol is 45%.

However, particularly preferred examples of solubilizing agents suitable for the topical composition with systemic effect according to the invention are:

- a ternary mixture 95° ethanol / water / propyleneglycol, in which the percentage of 95° ethanol
25 varies from 30 to 50%, that of water from 30 to 60%, and that of propyleneglycol from 2 to 20%; preferably, the percentage of 95° ethanol is 45%, that of water is 45%, and that of propyleneglycol is 8%,
- a quaternary mixture 95° ethanol / water / Labrasol® / propyleneglycol, in which the percentage of 95° ethanol varies from 30 to 50%, that of water from 30 to 60%, that of Labrasol® from 3 to
30 7% and that of propyleneglycol from 2 to 20%; preferably, the percentage of 95° ethanol is 45%, that of water is 33.5%, that of Labrasol® is 5% and that of propyleneglycol is 15%;

Of the substances regarded as absorption promoters, or "enhancers", the most used are derivatives of glycol, sulfoxides, surfactants, fatty acids and terpene derivatives.

As examples of absorption promoters, we may mention oleic acid, oleic alcohol, a triglyceride of decanoic and octanoic acids (for example, as marketed under the trade-name Miglyol 812®), isopropyl myristate, propyleneglycol dipelargonate, 2n-nonyl-1.3-dioxolane, octyl dodecyl myristate, isopropylidene glycerol (for example as marketed under the trade-name Solketal), α -tocopheryl propyleneglycol 1000 succinate (for example as marketed under the trade-name Vitamin E TPGS), monoethyl ether of diethyleneglycol (for example as marketed under the trade-name Transcutol®).

The absorption promoter that is more particularly suitable in the present invention is chosen from the group comprising isopropylidene glycerol, α -tocopheryl propyleneglycol 1000 succinate and monoethyl ether of diethyleneglycol.

However, the preferred absorption promoter is isopropylideneglycerol.

The forms envisaged for ensuring penetration of the active principle through the skin will be either gels, or occlusive gelled preparations.

The choice of gelling agents and film-forming agents is also important in the compositions according to the invention.

The gelling agents are substances which thicken and alter the viscosity of a liquid vehicle thus constituting a three-dimensional colloidal network, the gel.

There are various kinds of gelling agents: natural gelling agents (mineral, vegetable, animal), synthetic agents and semi-synthetic agents.

Examples of natural gelling agents are guar gum, extracts from algae (alginates, carrageenans, agar), polysaccharides (xanthan gum, gum arabic, tragacanth), starches, pectins, etc.

Examples of synthetic or semi-synthetic gelling agents are cellulose derivatives, especially those obtained by esterification or etherification of cellulose, and acrylic derivatives. The category of acrylic derivatives includes carbomers, polycarbophils, and acrylates.

In the present invention, the gelling agent is chosen from the group comprising cellulose derivatives and acrylic derivatives.

The cellulose derivatives include:

- methylcelluloses (Methocel, Metolose),
- ethylcelluloses (Ethocel, Aquacoat®)
- hydroxypropylmethylcelluloses (Kenal Methocel, Hypromellose),
- hydroxyethylcelluloses (Cellosize, Natrosol),
- hydroxypropylcelluloses (Klucel),
- carboxymethylcelluloses in the sodium or calcium form (Akucell, Nymcel Tylose CB),

The choice of a polymer is made from the Metolose range, from the company Shin Etsu. For each of these types, there are different degrees (or grades) depending on the substituents and the degree of substitution, which give different viscosities to the polymer solutions. There is a classification of the celluloses according to their adhesive potential. The choice of grade is important because the adhesive power of the cellulose derivative varies in relation to the latter.

According to the present invention, a particularly suitable cellulose derivative is hydroxypropylmethylcellulose, and especially hydroxypropylmethylcellulose of grade 60 SH 4000. In fact, grade 60 SH has the most suitable properties: good solubility in organic solvents and high resistance to electrolytes. It also makes it possible to obtain transparent gels.

Among the acrylic derivatives, the carbomers are particularly suitable according to the present invention, and especially those marketed under the trade-names Carbopol® or Synthalen®.

The carbomers give formulations that are stable over time, and endow the formulation with reproducible rheological properties on account of their synthetic nature.

The existence of different degrees or grades results from the difference in molecular weight, degree of crosslinking, nature of the molecular arrangements and the polymerization solvent.

Thus, among the various grades of carbomers, we may mention those marketed by the Goodrich company under the trade-names Carbopol 974 P ®, Carbopol 980 ®, Carbopol 1382 ® and Carbopol 2020 ®, or similar products such as the Synthalens from 3 V France, as they are (Synthalen K, L, M) or preneutralized, for example Synthalen PNC ®.

However, according to the present invention, the carbomers marketed under the trade-names Carbopol 980®, Carbopol 1382® and Synthalen K® are particularly suitable and offer considerable advantages, as they are fluidized in contact with the electrolytes of the skin and thus prevent deposition of polymer, which could hamper passage of the active principle.

The film-forming agents used are those that are employed for producing solutions for enrobing and coating, as most of them are obtained from the biomedical or food industry and are suitable for human application.

These film-forming agents can be placed in different groups according to their solubility.

Whatever the film-forming agent, the quality of the film-forming gel obtained or of the film-forming solution obtained depends on the percentage of film-forming agent, the type of solvent, the presence and nature of the plasticizer.

According to the present invention, the film-forming agent is chosen from the group comprising cellulose derivatives, methacrylic derivatives and polyvinylpyrrolidone derivatives.

Among the cellulose derivatives, we may mention:

- hydroxypropylmethylcellulose acetate succinate, and especially that marketed by the Seppic company under the trade-name Acoat AS-LF®,
- an aqueous dispersion of cellulose acetophthalate containing 70% water, 23% cellulose acetophthalate and 7% poloxamer, and especially that marketed by the Seppic company under the trade-name Aquacoat CPD®,
- an aqueous dispersion of ethylcellulose, cetyl alcohol and sodium lauryl sulphate, and especially that marketed by the Seppic company under the trade-name Aquacoat ECD 30®,
- ethylcellulose.

Among the methacrylic derivatives, we may mention:

- an aqueous dispersion of an anionic copolymer of methacrylic acid and ethyl acrylate (type C), especially that containing 30% of dry copolymer, 0.7% of sodium lauryl sulphate and 2.3% of Polysorbate 80 NF, and marketed under the trade-name Eudragit L30 D55® (Rohm & Haas),
- a copolymer of acrylic acid and methacrylic ester (type A), especially that marketed under the trade-name Eudragit RL 100® (Rohm & Haas).

Among the derivatives of polyvinylpyrrolidone, we may mention:

- a povidone, of formula $(C_6H_9NO)_n$ whose molecular weight is of the order of 360 000, marketed under the trade-name Kollidon 90®
- polyvinylpyrrolidone / vinylacetate 64 copolymer, of formula $(C_6H_9NO)_n \times (C_4H_6O_2)_m$ whose molecular weight is: $(111.1)_n \times (86.1)_m$.

- homopolymers of polyvinyl alcohol

In the present invention, the particularly suitable cellulose derivative is hydroxypropylmethylcellulose acetate succinate, the particularly suitable methacrylic derivative is an aqueous dispersion of an anionic copolymer of methacrylic acid and ethyl acrylate, and the particularly suitable derivative of polyvinylpyrrolidone is a povidone.

The topical hormonal compositions with systemic effect according to the invention can additionally contain other excipients that are complexing agents, neutralizing agents such as disodium edetate (EDTA), triethanolamine (TEA) and/or plasticizers such as diethyl phthalate and triacetin.

A particularly suitable topical hormonal composition according to the invention is a composition in the form of gel or film-forming gel, with a content of nomegestrol or nomegestrol acetate of 0.4 wt.% of the total composition, a pH between 6 and 7, and a viscosity between 1000 and 2000 mPas.

The method of preparation of the compositions with systemic effect according to the invention varies depending on the particular nature of the compositions that are to be produced, namely a gel, a film-forming gel or a film-forming solution.

• METHOD OF PREPARATION OF THE GELS

Similarly, during the preparation of compositions in the form of gel, the manner of preparation will not be exactly the same, depending on the type of gelling agent used. Thus, in the preparation of gels, with regard to the gelling agent a distinction is made between synthetic acrylic derivatives and cellulose derivatives.

- Preparation from acrylic derivatives

The important steps in the preparation of a gel are dispersion of the gelling agent in the solubilizing agent (this dispersion will largely determine the quality of the preparation obtained), stirring, hydration, swelling and finally gelling.

Dispersion and stirring: wetting

The acrylic derivative is suspended with stirring in the solvent (solubilizing agent). Stirring must be moderate, otherwise the acrylic polymer is degraded by shearing and loses its efficacy.

Hydration and swelling of the polymers

To prevent the formation of partially hydrated lumps, it is recommended to incorporate the polymers by sieving, to facilitate wetting and hydration of the powder and permit them to form a network. This step is promoted by wetting the powder beforehand, in the most polar solvent in the case of a solvent system.

Gelling: neutralization of the dispersion obtained

The pH of such a suspension is close to 3 (the pH is a function of the concentration of polymer, and therefore of carboxyl groups). Mineral bases such as sodium hydroxide, potassium hydroxide or ammonium hydroxide are used when the solvents in the formulation are aqueous, and organic bases such as amines (triethanolamine, tromethamine or TRIS etc.) when they are nonpolar or only slightly polar. Addition of these agents causes spontaneous thickening through formation of water-soluble salts of polymer resins.

An example of preparation of a gel whose gelling agent is an acrylic derivative is characterized in that:

- norgestrel acetate and EDTA are dissolved in the solvent system water / 95° ethanol / propyleneglycol with stirring at 300 rev/min (# 30 min);
- the acrylic polymer is dispersed in small portions in the solution of active principle with stirring at 100 rev/min;
- the acrylic polymer is left to swell for 2 hours with stirring at 200 rev/min;
- the dispersion is neutralized with triethanolamine dissolved in a portion of water taken from the quantity to be incorporated in the formulation; stirring is slowed to 100 rev/min during neutralization to avoid incorporation of air bubbles;
- it is stirred for 30 min at 150 rev/min to homogenize the gel obtained.

- Preparation from cellulose derivatives

Gels formulated on the basis of cellulose derivatives do not need to be neutralized, but it will sometimes be necessary to adjust their pH by means of organic amines or inorganic hydroxides, depending on the type of solvent in the formulation.

The viscosity obtained depends on the type and quantity of cellulose derivative used.

An example of preparation of a gel whose gelling agent is a cellulose derivative is characterized in that:

- norgestrel acetate and EDTA are dissolved in the solvent system water / 95° ethanol / propyleneglycol with stirring at 300 rev/min (# 30 min);

- the cellulose polymer is dispersed in small portions in the solution of active principle with stirring at 100 rev/min;
- the cellulose polymer is left to swell for 1 hour with stirring at 250 rev/min;
- the pH is adjusted, if necessary, with triethanolamine dissolved in water with stirring at 100 rev/min;
- it is stirred for 30 min at 150 rev/min to homogenize the gel obtained.

• FILM-FORMING GELS (or "FILMING GELS") AND FILM-FORMING SOLUTIONS (or "FILMING SOLUTIONS")

These forms are envisaged because, when they are applied to the skin, on drying they form a kind of occlusive film, which is sufficient to increase the hydration of the skin and create new sites for passage, thus improving the diffusion of the active principle that they contain. However, the form obtained must penetrate or dry quickly, leaving a pleasant, non-sticky feel.

The film-forming agents used in the present invention are generally those used for production of enrobing solutions for tablets.

• METHOD OF PREPARATION OF FILM-FORMING SOLUTIONS

As with gels, in the preparation of compositions in the form of a film-forming solution, the manner of preparation will vary depending on the nature of the film-forming agent used.

- Preparation from solid film-forming agent:

The steps in preparation are:

- Dissolving of a plasticizer and of the active principle in the solvent mixture: the mixture containing the plasticizer and the active principle must be stirred for a sufficient time to obtain a solution.
 - Dispersion and solution of the film-forming agent:
- Dispersion must be carried out in small portions, with vigorous stirring. Stirring is continued until the film-forming agent has dissolved completely. Neutralization of the film-forming solution is carried out, if necessary, at the end of production, with slower stirring.

An example of preparation of a film-forming solution whose film-forming agent is solid is characterized in that:

- the quantities of ethanol, water and propyleneglycol required for the formulation are stirred at 250 rev/min for 10 min;

- 5 - EDTA and nomegestrol acetate are dissolved in the mixture obtained;
 - the plasticizer is added and stirring is continued for 30 min at 250 rev/min;
 - the film-forming agent is dispersed in small portions, stirring at the same speed, until it has dissolved completely; stirring is then continued for 1 hour;
 - the pH is adjusted by means of a solution of triethanolamine dissolved in a small amount of
- 10 water, taken from the quantity of water to be incorporated in the formulation, reducing stirring to 100 rev/min; the solution obtained is homogenized for 30 min.

- Preparation from a film-forming agent in aqueous dispersion

The steps in preparation are:

- 15 - Solution and plasticizing of the film-forming agent
- Incorporation of the mixture containing the active principle and the other excipients in small portions, with vigorous stirring

Neutralization is carried out at the end of production, with slower stirring.

20 An example of preparation of a film-forming solution whose film-forming agent is an aqueous dispersion is characterized in that:

- the water and a plasticizer are mixed at 250 rev/min; stirring for 30 min;
 - the dispersion of film-forming agent is added in small portions, stirring at 250 rev/min, until a homogeneous solution of the dispersion is obtained; stirring is continued for 1 hour;
- 25 - separately, EDTA and nomegestrol acetate are dissolved in the ethanol / propyleneglycol mixture; stirring is continued until they are dissolved completely;
- the alcoholic solution of active principle is added in small portions to the aqueous solution, with stirring at 250 rev/min; the solution obtained is stirred for 1 hour to homogenize it;
 - the solution is neutralized with triethanolamine dissolved in water, with slower stirring; the
- 30 solution obtained is homogenized for 30 min.

• METHOD OF PREPARATION OF FILM-FORMING GELS OR GELLED FILMS

The film-forming gels are obtained by gelling of film-forming solutions.

First, two solutions are prepared separately:

- an aqueous solution containing a dissolved plasticizer, in which the film-forming agent is dissolved completely with vigorous stirring;
- an alcoholic solution containing the other excipients of the formulation and in which the active principle is dissolved; the gelling agent is dispersed in it and left to swell.

Then the alcoholic solution is mixed into the aqueous solution and the solution is gelled with triethanolamine.

An example of preparation of a film-forming gel is characterized in that:

- the plasticizer is dissolved in water; stirring is carried out for 30 min at 250 rev/min;
- the film-forming agent is dispersed in it and stirred at 250 rev/min until it has dissolved completely (in the case of a solid film-forming agent) or until the dispersion is homogeneous; stirring is continued for 1 hour;
- separately, the EDTA and nomegestrol acetate are dissolved in the mixture of propyleneglycol and ethanol; the chosen gelling agent is dispersed and is left to swell for 2 hours, stirring at 150 rev/min;
- the alcoholic solution is mixed with the aqueous solution, and stirred for 1 hour at 150 rev/min;
- neutralization is carried out with triethanolamine dissolved in water, lowering the speed of stirring to 100 rev/min; the gel obtained is homogenized by stirring for 30 min.

METHOD OF EVALUATING PASSAGE OF THE ACTIVE PRINCIPLE THROUGH THE SKIN

The efficacy of the topical composition according to the invention is evaluated by demonstrating that the active principle it contains diffuses through the skin and is absorbed into the microcirculation in sufficient quantity to achieve the desired therapeutic effect.

In the present invention, passage of nomegestrol acetate through the skin is evaluated by measuring the radioactivity, using a molecule labelled with carbon 14. The method of evaluation of passage of the active principle using radiolabelled products makes it possible to detect low levels of the active principle, which represents a considerable advantage, bearing in mind the small quantities that diffuse through the skin.




The skin used in the various tests for evaluating percutaneous passage of the active principle was obtained from abdominal plastic surgery on female subjects in the age range from 40 to 45 years. Excess adipose tissue is removed from the skin, which is then cleaned and stored in a freezer at -70°C.




The topical compositions according to the invention are intended to be applied mainly to the skin of the abdomen, arms, thighs, etc.

EXPERIMENTAL SECTION

• EXAMPLE I

Fig. 1 illustrates passage of the active principle nomegestrol acetate (NAc) through the skin as a function of different quantities of nomegestrol acetate in the compositions according to the invention.

The symbols , , and  in Fig. 1 represent:

 Gel 0.11 % of NAc	 Gel 0.4 % of NAc	 Gel 0.8 % of NAc
--	---	---

These compositions are in the form of gel and their formulations are given in Table 1 below :

Table 1

FORM S	GELS		
FORMULATIONS (in %)			
Nomegestrol Acetate	0.40	0.80	0.11
Propyleneglycol	6.00	6.00	3.00
Transcutol®	5.00	5.00	-
Carbopol 1342®	0.50	0.50	-
Carbopol 940®	/	/	0.75
EDTA	0.05	0.05	0.05
Triethanolamine (TEA)	0.30	0.30	0.30
Demineralized water	42.75	42.35	45.79
Ethanol	45.00	45.00	50.00
pH (at 1 %)	6.7	6.5	6.7
NOMEGESTROL ACETATE (mg/g)	0.41	0.4	0.403

Passage of the active principle through the skin is evaluated by measuring the amount of active principle accumulated as a function of time. The amount of active principle accumulated

represents the total diffusion of the active principle through the skin over a specified period (24 or 48 h). In this example, it is expressed in ng.

Fig. 1 clearly shows that the poorest diffusion results are obtained with the gel with 0.11 % of NAc.

This gel at 0.11 % was tested in preliminary clinical trials : cf. example IV. Thus, it was established that this gel, despite its poorer results, was still able to achieve a systemic passage effect.

• EXAMPLE II

Investigation of the solubility of nomegestrol acetate (NAc)

1) - a) In a 95° ethanol / water binary mixture

The most effective system of solvent in aqueous-alcoholic mixture is determined.

Table 2 : Solubility of nomegestrol acetate as a function of the percentage of 95° ethanol

% 95° ethanol	Solubility of nomegestrol acetate in mg/ml
0	0.056
10	0.070
20	0.113
30	0.683
40	2.820
50	7.330
60	17.850
70	24.850
80	29.500
90	26.600
100	32.850

In aqueous-alcoholic mixture, solubility increases with the percentage of alcohol. The solubility profile shows that it is fairly low up to 40 % alcohol, then increases sharply between 40 and 80 %. Now, the percentage of alcohol permitted for forms for topical application is limited. Within these limits, the most effective solvent system for solubilizing nomegestrol acetate is between 40 and 60 % of alcohol.

- b) In a ternary mixture 95° ethanol / water / propyleneglycol

The effect of a ternary mixture of solvents, ethanol / water (45:55) / propyleneglycol on the solubility of nomegestrol acetate was determined.

- 5 We also examined the possibility of lowering the proportion of alcohol in the solvent, by means of this ternary mixture, while maintaining similar solubility ; for this we choose the influence of propyleneglycol on solubility in ethanol / water systems (40:60 and 30:70).

Table 3 : Solubility of nomegestrol acetate in various systems containing propyleneglycol (PG)

10

% Propyleneglycol	Solubilities (mg/ml)		
	System I	System II	System III
0	0.6	2.9	5.1
2	0.6	2.6	5.1
4	0.5	2.6	5.1
6	0.7	3.0	5.1
8	1.0	3.2	7.7
12	1.1	3.4	7.7
20	1.5	3.9	7.9

System I : $\left[\begin{array}{l} \text{Ethanol } 95^\circ : 30 \% \\ \text{Demineralized water} : 70 \% \end{array} \right.$

System II : $\left[\begin{array}{l} \text{Ethanol } 95^\circ : 40 \% \\ \text{Demineralized water} : 60 \% \end{array} \right.$

15 System III : $\left[\begin{array}{l} \text{Ethanol } 95^\circ : 45 \% \\ \text{Demineralized water} : 55 \% \end{array} \right.$

Table 3 is illustrated by Fig. 2.

The symbols \boxtimes , \blacklozenge , and π in Fig. 2 represent :

20

\boxtimes Solubility in system I	π Solubility in system III
\blacklozenge Solubility in system II	

25 In water / ethanol / propyleneglycol ternary mixture, the solubility of the active principle is improved with proportions of 8 % of propyleneglycol for a mixture with 45 % alcohol. It is for this system that we obtain the best solubility of the active principle. Propyleneglycol acts in synergy with the alcohol, on the solubility of nomegestrol acetate.

- c) In 95° ethanol / water / Labrasol / propyleneglycol mixture

Table 4 : Solubility of nomegestrol acetate in a system containing propyleneglycol

% Propyleneglycol	Solubility of l'nomegestrol acetate (mg/ml)
0	9.4
10	9.5
15	10.2

System : { Ethanol 95° : 45 %
Water : 50 %
Labrasol® : 5 %

Using Labrasol® alone at 5 % without propyleneglycol, the solubility increases in comparison with the results obtained with propyleneglycol. This solubility is increased further by combining propyleneglycol and Labrasol®.

2) - a) In a 95° ethanol / water / propyleneglycol / Solketal mixture

Table 5 : Solubility of nomegestrol acetate in an aqueous-alcoholic mixture containing propyleneglycol and/or Solketal

% Propyleneglycol	% Solketal	Solubility of nomegestrol acetate (mg/ml)
0	3	6.7
0	8	8.6
8	0	7.7
8	3	10.7

The solubility of nomegestrol acetate in the aqueous-alcoholic solvent mixture in the presence of 8 % of Solketal is greater than that obtained in the presence of 8 % of propyleneglycol alone. Combination of the two substances propyleneglycol and Solketal greatly increases solubility in the aqueous-alcoholic solvent mixture, in the proportion of 8 % of propyleneglycol / 3% of Solketal.

- b) In a 95° ethanol / water / propyleneglycol / vitamin E TPGS mixture

Table 6 : Solubility of nomegestrol acetate in an aqueous-alcoholic mixture containing propyleneglycol and/or vitamin E TPGS

% Propyleneglycol	% Vitamin E TPGS	Solubility of nomegestrol acetate (mg/ml)
0	3	7.65
0	8	11.10
8	0	7.70
8	3	12.50

The solubility of nomegestrol acetate is improved in the presence of vitamin E TPGS alone, relative to propyleneglycol, for a same proportion of 8 %. Incorporated at 3 %, it already gives results equivalent to propyleneglycol used at 8 %.

However, even better solubility is obtained when these two substances are combined in the proportion of 8 % of propyleneglycol / 3 % of vitamin E TPGS.

- c) In a 95° ethanol / water / propyleneglycol / Transcutol® mixture

Table 7 : Solubility of nomegestrol acetate in an aqueous-alcoholic mixture containing propyleneglycol and/or Transcutol®

% Propyleneglycol	% Transcutol ®	Solubility of nomegestrol acetate (mg/ml)
8	0	7.70
8	3	7.95
0	8	10.70
3	8	10.60

The solubility of nomegestrol acetate is improved in the presence of 8 % of Transcutol® alone, relative to propyleneglycol at the same proportion. On combining these two substances, the same solubility is obtained using the proportions 8 % Transcutol / 3 % propyleneglycol.

Reversing the proportions does not improve the solubility of the active principle in comparison with that obtained in propyleneglycol alone.

Conclusion:

Aqueous-alcoholic mixtures containing :

- 8 % of propyleneglycol and 3 % of Solketal,
- or 8 % of propyleneglycol and 3 % of vitamin E TPGS,
- or 3 % of propyleneglycol and 8 % of Transcutol®,

are particularly suitable for good solubility of the active principle.

• EXAMPLE III

1/ Investigation of formulations in the form of gels

Among the substances chosen for their qualities as absorption promoters, Solketal and vitamin E TPGS are particularly suitable as they also prove capable of improving the solubility of nomegestrol acetate aqueous-alcoholic mixture and propyleneglycol. The promoting action of the three promoters was investigated by incorporating them in formulations containing an aqueous-alcoholic gel at 45 % alcohol and including 8 % of propyleneglycol and 3 % of the promoter.

These formulations in the form of gels were tested for passage through the skin.

The gels used meet the requirements for pH, viscosity, concentration and appearance.

The four gels investigated were designated « G36-264, G36-276, G32-104 and G37-113 » and their formulations are presented in Table 8 below :

Table 8 :

FORM S	G E L S			
	G36-264	G36-276	G32-104 reference	G37-113 reference
References				
Formulations in %				
Nomegestrol Acetate	0.4	0.4	0.4	0.4
Propyleneglycol	8	8	8	8
Transcutol®	/	/	3	3
Solketal	3	/	/	/
Vit E TPGS	/	3	/	/
HPMC 60SH4000	/	/	/	/
Carbopol 1382®	/	/	0.5	/
Carbopol 980®	0.5	0.5	/	0.6
EDTA disodium edetate	0.05	0.05	0.05	0.05
TEA	0.4	0.3	0.3	0.25
Kollidon 90®	/	/	/	/
Aquat AS-LF®	/	/	/	/
Eudragit L30D55®	/	/	/	/
Diethyl phthalate	/	/	/	/
Ethanol 95	45	45	45	45
Demineralized water	42.65	42.75	42.75	42.85
pH solution at 1 %	6.9	6.92	6.6	6.37
Viscosity mPa.s	1150	1020	1400	1400
NAC content (%)	0.41	0.4	0.403	0.393

The main differences in the composition of these gels relate firstly to the choice of absorption promoter ("enhancer") and secondly to the choice of gelling agent, which is either Carbopol 980® or Carbopol 1382®.

Passage of the active principle through the skin is evaluated by measuring:

- the cumulative amount of active principle as a function of time (cf. example I),
- the cumulative percentage of active principle as a function of time,
- and the rate of diffusion of active principle as a function of time.

The cumulative percentage of active principle is the total percentage of diffusion of the active principle through the skin in a given time interval.

The rate of diffusion of the active principle is expressed in $\mu\text{g} / \text{cm}^2 / \text{h}$: it can be used for determining the kinetics of diffusion of the active principle over time.

The method used for evaluating passage of the active principle also makes it possible to determine the distribution of nomegestrol acetate in the various structures of the skin after diffusion.

- 5 Table 9 shows the cumulative percentage of nomegestrol acetate as a function of time, as well as (cf. last 3 lines) the sites of distribution of the active principle in the skin structures, i.e. the content of nomegestrol acetate in the various layers (epidermis + dermis) of the skin.

Table 9 :

Time, h	cumulative % in μg			
	G E L S			
	G37-113	G32-104	G36-276	G36-264
	8 % PG 3 % Trans 0.6 % C980	8 % PG 3 % Trans 0.5 % C1382	8 % PG 3 % Vit E 0.5 % C980	8 % PG 3 % Solketal 0.5 % C980
0	0	0	0	0
2	0.042	0.058	0.105	0.183
4	0.088	0.135	0.193	0.371
6	0.137	0.227	0.275	0.554
8	0.19	0.329	0.347	0.732
10	0.239	0.402	0.405	0.894
24	0.575	1.117	0.667	1.926
Epidermis	10.05	14.82	8.15	16.82
Dermis	4.33	6.97	4.68	4.84
Washing	67.63	61.15	72.69	60.35

10 Table 9 is illustrated by Figs. 3 and 4.

Fig. 3 shows the influence of enhancer (promoter) and of Carbopol® on passage of the systemic gel of nomegestrol acetate through the skin.

The following promoters were compared : Transcutol® (Tr), Solketal (Sol) and Vitamin E TPGS (Vit E).

15 The symbols \square , π , \blacksquare and \boxtimes in Fig. 3 represent :

\square G37-113 (3 % Tr)	\blacksquare G32-104 (3 % Tr) - Ref.
π G36-276 (3 % Vit E)	\boxtimes G36-264 (3 % Sol)




Fig. 4 shows the distribution of nomegestrol acetate in the structures of the skin.

The symbols , , , and  in Fig. 4 represent :

 G37-113 (3 % Tr)	 G32-104 (3 % Tr) - Ref.
 G36-276 (3 % Vit E)	 G36-264 (3 % Sol)

From the values of cumulative percentage of active principle, it is possible to deduce the values of cumulative amount and rate of diffusion.

Fig. 5 shows the rate of diffusion of nomegestrol acetate as a function of time.

The symbols , π ,  and  in Fig. 5 have the same meaning as in Fig. 3.

The kinetics of diffusion of nomegestrol acetate with the compositions in the form of gels is of the patch type, with constant diffusion.

Conclusion:

Solketal gives a greater improvement of passage of nomegestrol acetate through the skin than Vitamin E and Transcutol® if we compare the results obtained with those of the G32-104 reference gel.

Thus, better diffusion is obtained using Solketal rather than Vitamin E TPGS, whereas the solubility of the active principle is better for the latter (cf. Tables 5 and 6).

The same comment can be made regarding an aqueous-alcoholic system with the propyleneglycol / Transcutol® mixture: if we consider the four possible combinations 8:0 - 8:3 - 3:8 and 0:8, the diffusion obtained is best for the 8:3 mixture. However, solubility is poorest for the latter (cf. Table 7).

The active principle must have some affinity for the solvent if it is to be dissolved completely. However, it must not be too great, so that the partition coefficient between the vehicle and the skin is oriented in favour of diffusion through the skin.

Diffusion tests in static flow with radiolabelled active principle were carried out for gels containing the two types of absorption promoters adopted, versus two reference gels: gel G32-104 (a grade of carbopol different from G37-113 and the other two), for which the best diffusion has been obtained, and gel G37-113, of identical composition to the two gels tested (same grade of Carbopol®) containing Transcutol®.

Examining the action of the absorption promoters adopted, on passage of labelled nomegestrol acetate through the skin, we observe a clear increase in diffusion in the presence of Solketal relative to the G32-104 reference gel containing Transcutol®.

- 5 Vitamin E TPGS, used in the same conditions, does not improve passage compared with gel G32-104. On the other hand, if we consider gel G37-113, the diffusion obtained is slightly better with vitamin E, and is increased markedly with Solketal.

When we examine the quantitative distribution of the active principle (cf. Fig. 4) in the structures
10 of the skin, for the active principle of gel G36-264 and of the reference gel G32-104 we find similar concentrations in the epidermis and in the dermis. There is a lower level in the epidermis for gels G37-113 and G36-276.

The above tests also confirm that there is a difference in diffusion of active principle depending
15 on the grade of carbomer used in the formulation (the formulations of gels G37-113 and G32-104 are quantitatively and qualitatively identical apart from this). It appears that diffusion is better in the presence of Carbopol 1382 ®, just as with distribution in the structures of the skin.

Examining the results obtained with respect to adhesion, in tests carried out on these gels G36-
20 264 and G36-276, it is found that the adhesive character of the gel containing Solketal is slightly better than that of the gel containing vitamin E. Note that these two gels contain the same proportions of the excipients, apart from the type of promoter.

Conclusion

A topical hormonal composition with systemic effect currently preferred according to the present
25 invention is a composition in the form of gel containing :

- 0.4 % nomegestrol acetate
- 8 % propyleneglycol
- 3 % Solketal
- 0.5 % Carbopol 980 or 1382®
- 30 - 45 % ethanol 95°
- 0.05 % EDTA, 0.4 % TEA and demineralized water to give 100%.

2 / Investigations of formulations in the form of film-forming solutions

The 5 film-forming solutions investigated were designated « G36-259, G36-261, G36-263, G36-266 and G36-277 » and their formulations are shown in Table 10 below :

Table 10

<i>FORMS</i>	<i>FILM-FORMING SOLUTIONS</i>				
References	G36-259	G36-261	G36-263	G36-266	G36-277
Formulations in %					
Nomegestrol Acetate	0.4	0.4	0.4	0.4	0.4
Propyleneglycol	8	8	8	8	8
Transcutol®	/	/	/	/	/
Solketal	/	/	/	3	/
Vit E TPGS	/	/	/	/	3
HPMC 60SH4000	/	/	/	/	/
Carbopol 1382®	/	/	/	/	/
Carbopol 980®	/	/	/	/	/
EDTA	0.05	0.05	0.05	0.05	0.05
TEA	/	0.8	0.3	0.05	0.05
Kollidon 90®	5	/	/	5	5
Aquat AS-LF®	/	10	/	/	/
Eudragit L30D55®	/	/	10	/	/
Diethyl phthalate	/	3	2	/	/
Ethanol 95	43.35	40	40	43.25	43.25
Demineralized water	43.2	37.75	39.25	40.25	40.25
pH solution at 1 %	6.25	6.16	6.24	6.83	6.34
Viscosity MPa.s					
NAC content (%)	0.41	0.42	0.43	0.40	0.40

The main differences in the composition of these film-forming solutions relate to the choice of film-forming agent and the choice of addition, or not, of an absorption promoter or of a plasticizer.

Tests were carried out for the film-forming solutions in comparison with gel G32-104 as reference.

Table 11 shows the cumulative percentage of nomegestrol acetate as a function of time and the distribution of nomegestrol acetate in the cutaneous structures.

Table 11

Time, h	Cumulative percentage					
	GELS	FILM-FORMING SOLUTIONS				
	G32-104	G36-261	G36-263	G36-259	G36-266	G36-277
	8% PG 3% Trans 0.5% C1382	8% PG 10% Aqoat	8 % PG 10% Eudr	8 % PG 5% Koll	8% PG 3% Solk 5% Koll	8% PG 3% TPGS 5% Koll
0	0	0	0	0	0	0
2	0.062	0.088	0.092	0.077	0.067	0.064
4	0.123	0.153	0.152	0.14	0.116	0.112
6	0.185	0.206	0.197	0.203	0.165	0.153
8	0.252	0.251	0.239	0.269	0.211	0.193
10	0.342	0.3	0.289	0.349	0.269	0.242
24	0.799	0.487	0.515	0.699	0.539	0.474
Epidermis	8.24	5.69	2.78	9.14	6.41	4.43
Dermis	5.56	1.93	2.26	4.58	3.2	4.76
Washing	72.18	97.96	98.85	94.33	91.37	95.24

Table 11 is illustrated by Figs. 6 and 7.

Fig. 6 shows the influence of the absorption promoter and of the film-forming agent on passage of the systemic film of nomegestrol acetate through the skin.

The symbols □, ■, ♪, ★, π and △ in Fig. 6 represent :

□	G32-104 (3%Tr) -Ref.-	♪	G36-261 (10%Aqoat)
■	G36-259 (5%Kol)	★	G36-266 (3% Solk/5%Kol)
		π	G36-263 (10%Eudrag)
		△	G36-277 (3%TPGS/5%Kol)







- 15 The amounts of active principle that had diffused from these forms are less in all cases than the amounts obtained by application of the non-filming gel G32-104, regardless of the polymer considered.

It can be seen that the solution containing only Kollidon® gives diffusion that is closest to that of the reference gel. The other two polymers produce similar diffusions.

- 20 Solutions combining Kollidon® and a promoter, such as Solketal or vitamin E TPGS, give poorer diffusions of active principle compared with solution G36-259 without promoter.

Fig. 7 shows the distribution of nomegestrol acetate in the skin structures.

The symbols , , , ,  and  in Fig. 7 represent:

	G32-104 (3%Tr) -Ref.-		G36-261 (10%Aquat)
	G36-259 (5%Kol)		G36-266 (3% Solk/5%Kol)
		G36-263 (10%Eudrag)	
		G36-277 (3%TPGS/5%Kol)	

It can be seen that distribution is best for Kollidon®. It is equivalent to that found for the reference gel. The results obtained for the solutions of Aquat® and of Eudragit® remain low.

The diffusion results obtained with the film-forming solutions are slightly better than those obtained with the gel with 0.11 % of nomegestrol acetate (cf. Table 1, example I). However, it should be noted that with regard to Eudragit® and Aquat®, the solutions prepared only contain propyleneglycol, without any other promoter, in contrast to the reference gel.

Conclusion :

A topical hormonal composition with systemic effect according to the present invention will be, for example, a composition in the form of film-forming solution containing :

- 0.4 % of nomegestrol acetate
- 8 % of propyleneglycol
- 5 % of Kollidon 90®
- 43.35 % of 95° ethanol
- 0.05 % of EDTA and demineralized water to give 100%.

3 / Investigation of formulations in the form of film-forming gels or gelled films

The three film-forming gels investigated were designated « G36-260, G36-262 and G36-267 » and their formulations are shown in Table 12 below.

Table 12

FORMS	GELS		
	Film-forming	G36-262	G36-267
References	G36-260		
Formulations in %			
Nomegestrol Acetate	0.4	0.4	0.4
Propyleneglycol	8	8	8
Transcutol®	/	/	/
Solketal	/	/	/
Vit E TPGS	/	/	/
HPMC 60SH4000	/	/	1
Carbopol 1382®	/	/	/
Carbopol 980®	0.5	0.75	/
EDTA	0.05	0.05	0.05
TEA	0.1	0.9	0.4
Kollidon 90®	5	/	/
Aquat AS-LF®	/	10	/
Eudragit L30D55®	/	/	10
Diethyl phthalate	/	3	2
Ethanol 95	43	40	40
Demineralized water	42.95	36.9	38.15
pH solution at 1 %	6.36	6.2	6.17
Viscosity mPa.s	1750	1050	1150
NAC content (%)	0.41	0.405	0.40

The main differences in the composition of these film-forming gels relate to the choice of gelling agent and of film-forming agent.

These tests were carried out for the film-forming gels in comparison with gels G32-104 and G37-113 as reference.

Table 13 shows the cumulative percentage of nomegestrol acetate as a function of time and the distribution of nomegestrol acetate in the structures of the skin.

Time h	% Cumulative				
	GEL		FILM-FORMING GEL		
	G32-104	G37-113	G36-260	G36-262	G36-267
	8% PG 3% Trans 0.5% C1382	8% PG 3% Trans 0.6% C980	8% PG 5% Kollidon 0.5% C980	8 % PG 10% aqoat 0.75% C980	8 % PG 10% Eudr 1% HPMC
0	0.000	0.000	0.000	0.000	0.000
2	0.058	0.024	0.121	0.251	0.251
4	0.135	0.088	0.206	0.414	0.392
6	0.227	0.137	0.278	0.533	0.506
8	0.329	0.190	0.334	0.626	0.590
10	0.402	0.239	0.379	0.694	0.650
24	1.117	0.575	0.598	0.994	0.913
Epidermis	14.82	10.05	11.42	16.36	11.21
Dermis	6.97	4.33	4.81	1.53	1.39
Washing	61.15	67.63	65.83	72.35	90.42

Table 13 is illustrated by Figs. 8 and 9.

Fig. 8 shows the influence of the film-forming agent on passage of the systemic film-forming gel of nomegestrol acetate through the skin.

The symbols \triangle , \boxtimes , \blacksquare , \spadesuit and π in Fig. 8 represent :

\triangle G32-104 (C1382)	\blacksquare G36-260 (Kol/C980)	π G36-262 (aq/C980)
\boxtimes G36-267 (Eud/HPMC)	\spadesuit G37-113 (C980)	

Examining all of the polymers, the diffusion of nomegestrol acetate from film-forming gels of Aqoat® and of Eudragit® is better than for the G32-104 « reference » gel up to 10 hours. Beyond that, the trend is slightly reversed. If we consider the non-filming gel 113, containing a different carbopol® from that in gel G32-104, the results obtained for all the film-forming gels are better, regardless of the polymer considered.

Note that diffusion of the active principle is similar for Aqoat® and Eudragit®.

On the other hand, it is much lower for Kollidon®.

Fig. 9 shows the distribution of the active principle in the structures of the skin.

The symbols ▨ , ▩ , ▧ , ▦ , ▥ in Fig. 9 represent :

▨ G32-104 (C1382)	▩ G36-260 (Kol/C980)	▧ G36-262 (aq/C980)
▦ G36-267 (Eud/HPMC)	▥ G37-113 (C980)	

It can be seen that the distribution varies from one polymer to another : relative to the G32-104 reference gel, the distribution in the epidermis is similar for Acoat®, but lower for Kollidon® and Eudragit®. The distribution in the dermis is lower for Acoat® and Eudragit®, but higher for Kollidon®.

Conclusion :

Examples of topical hormonal compositions with systemic effect according to the invention are, for example, compositions in the form of film-forming gel containing :

- 0.4 % of nomegestrol acetate
- 8 % of propyleneglycol
- 0.75 % of Carbopol 980®
- 10 % of Acoat AS-LF®
- 40 % of ethanol 95°
- 3 % of diethyl phthalate, 0.05 % of EDTA, 0.9 % of TEA and demineralized water to 100%,

or, compositions in the form of film-forming gel containing :

- 0.4 % of nomegestrol acetate
- 8 % of propyleneglycol
- 1 % of HPMC 60 SH 4000
- 10 % of Eudragit L 30 D 55®
- 40 % of ethanol 95°
- 2 % of diethyl phthalate, 0.05 % of EDTA, 0.4 % of TEA and demineralized water to 100%.





4 / Comparison between film-forming solutions and film-forming gels

Table 14 shows the cumulative percentage of nomegestrol acetate as a function of time, and the distribution of nomegestrol acetate in the structures of the skin.

Time h	% cumulative					
	FILM-FORMING SOLUTIONS			FILM-FORMING GELS		
	G36-259	G36-261	G36-263	G36-260	G36-262	G36-267
	8% PG 5 % Koll	8% PG 10% Aqoat	8 % PG 10% Eudr	8 % PG 5% Koll 0.5%C980	8% PG 10%Aqoat 0.75%C980	8% PG 10%Eudr 1%HPMC
0	0	0	0	0	0	0
2	0.077	0.088	0.092	0.121	0.251	0.251
4	0.14	0.153	0.152	0.206	0.414	0.392
6	0.203	0.206	0.197	0.276	0.533	0.506
8	0.269	0.251	0.239	0.334	0.626	0.59
10	0.349	0.3	0.289	0.379	0.694	0.65
24	0.699	0.487	0.515	0.598	0.994	0.913
Epidermis	9.14	5.69	2.78	11.42	16.36	11.21
Dermis	4.58	1.93	2.26	4.81	1.53	1.39
Washing	94.33	97.96	98.85	65.83	72.35	90.42

Table 14 is illustrated by Figs. 10 and 11.

Fig. 10 makes it possible to compare film-forming solutions and film-forming gels of nomegestrol acetate with systemic effects.

The symbols , , , , π and ρ in Fig. 10 represent :











<u>Film-forming solutions</u>		
 G36-259 (5% Kollidon)	 G36-261 (10 % Aqoat)	π G36-263 (10 % Eudrag)
<u>Film-forming gels</u>		
 G36-260 (5% Kol/C980)	 G36-262 (10%Aqoat/C980)	ρ G36-267 (10 % Eudr/HPMC)

Fig. 11 shows the distribution of the active principle in the structures of the skin.

The symbols , ,  (column 2, fig. 11), ,  (column 3, fig. 11) and  (last column) in Fig. 11 represent :







 G36-259 (5% Kollidon)	 G36-261 (10 % Aqoat)	 G36-263 (10 % Eudrag)
 G36-260 (5% Kol/C980)	 G36-262 (10%Aqoat/C980)	 G36-267 (10 % Eudr/HPMC)

Fig. 12 compares the diffusion of nomegestrol acetate with compositions in the form of film-forming gel and with compositions in the form of film-forming solution.

The symbols in Fig. 12 have the same meanings as in Fig. 10.

In contrast to Fig. 5 (diffusion flux with compositions in the form of gel), the kinetics of diffusion is not kinetics with constant diffusion, but very quickly (after 2 hours) displays maximum diffusion which then decreases quite quickly. This is especially true of the film-forming gels G36-262 and G36-267. Thus, two types of flux can be distinguished : flux with more or less constant diffusion, and other forms which very quickly exhibit a maximum diffusion peak.

Thus, the film-forming gels are better suited than a film-forming solution for optimization of percutaneous distribution of nomegestrol acetate. More particularly, the mere presence of a cellulosic (Aqoat®, G36-262) or acrylic (Eudragit®, G36-267) film-forming agent in a film-forming gel makes it possible to achieve good diffusion of the active principle. The film that forms, in both cases, is stronger, more cohesive, and seems to permit release of the active principle.

Accordingly, it is possible that topical hormonal compositions with systemic effect in the form of film-forming gel can be combined with 3 % of Solketal to obtain a synergistic action and further improve the diffusion of nomegestrol acetate.

4 / Conclusion

Film-forming or filming solutions generally only permit diffusion of active principle less than that obtained for the reference gel (G32-104).

On the other hand, film-forming gels of Aqoat® (G36-262) and of Eudragit® (G36-267) can provide considerable diffusion of active principle, if we consider the formulations that do not contain any absorption promoter.

Solketal is an absorption promoter that seems to have an action on the diffusion of nomegestrol acetate through the skin ; thus, in an aqueous-alcoholic system and when combined with propyleneglycol in the proportions (3:8), it greatly improves its solubility in the vehicle and its passage through the skin.

Thus, a particularly suitable example of a topical hormonal composition with systemic effect according to the invention is a composition that is in the form of a gel or a film-forming gel and

contains, in an aqueous-alcoholic mixture, 8 % of propyleneglycol and 3 % of isopropylidene glycerol.

• **EXAMPLE IV**

Preliminary clinical trials

5 In these examples, clinical trials were carried out on women, with the gel containing 0.11 % of nomegestrol acetate, whose formulation is given in Table 1 of example I.

1/ Clinical example No. 1

Twentry-four volunteers, women in good health and in the period of ovarian activity, with
10 average age of 23.5 years, were treated on 15 consecutive days with 4 mg of nomegestrol acetate in a gel applied every day to both breasts.

Repeated blood samples were taken in the hours following the first and last administration, on 9
occasions (6 times before application of the gel and 3 times 3 hours afterwards), between the 2nd
15 and the 14th day of treatment. The plasma of these samples was analysed for nomegestrol acetate by liquid chromatography combined with mass spectrometry.

From the very first day of treatment, nomegestrol acetate was detected in all the subjects. The
maximum concentration was evaluated at 0.25 ± 0.027 ng/ml and the area under the curve from 0
20 to 48 hours, of 6.08 ± 0.775 ng/ml per h, the mean values forming a plateau between 0.10 and 0.17 ng/ml.

After the last administration, the maximum concentration was 0.65 ± 0.073 ng/ml and the area
under the curve from 0 to 48 hours was 18.43 ± 2.091 ng/ml per h, nomegestrol acetate still
25 being detected in the plasma 72 hours after the last application (at a level of 0.19 ± 0.027 ng/ml).

A state of equilibrium is obtained after the 3rd day of treatment. We then observe mean values
that remain on a plateau, with little variation, between 0.42 and 0.65 ng/ml.

30 2/ Clinical example No. 2

Six menopausal women, with ages ranging from 56 to 66 years and without hormone
replacement treatment for 2 months, were monitored for 2 consecutive cycles of 25 days,
separated by a therapeutic window of 6 days.

During each day, they were given one tablet of oral oestradiol per day and, during the 15 days of the second cycle, 4 mg of nomegestrol acetate applied as a gel on the abdominal skin. At the end of each cycle, the plasma was analysed for nomegestrol acetate, the occurrence of genital bleeding was noted and a biopsy of the endometrium was carried out.

In contrast to what was observed in the first cycle (oestradiol alone), during the second cycle, administration of the gel of nomegestrol acetate showed that the progestogen could be detected in the plasma at levels between 0.39 and 0.76 ng/ml (0.62 ng/ml on average) and that these levels were sufficient to cause secretory transformation of the endometrium and produce genital bleeding, on average 5 days after the end of the second cycle.

3/ Clinical example No. 3

One hundred and thirteen premenopausal women, who had been suffering from breast pains for more than 3 months and for at least 7 days per cycle, were treated for an average time of 130 days with 4 mg of nomegestrol acetate applied each day, on the last 15 days of the menstrual cycle, in the form of a gel on both breasts.

Efficacy was assessed after 3 months and at the end of treatment using a visual analogue scale for quantifying breast pain.

This evaluation demonstrated that the nomegestrol acetate gel led to a statistically significant decrease in the intensity and duration of the cycle of breast pain, starting from the 3rd month of treatment. After 6 cycles of treatment, the intensity had decreased by 48 % and the duration by 41 %.

In the course of this study, 55 women were analysed for nomegestrol acetate in the blood, and values of 0.44 ± 0.30 (m \pm sd) ng/ml were obtained.

CLAIMS

1. Topical hormonal composition with systemic effect for the correction of progesterone deficiency in premenopausal women and for hormone replacement in menopausal women,
5 characterized in that it includes :

- as active principle, a progestogen derived from 19-nor progesterone,
- a vehicle permitting the systemic passage of the said active principle chosen from the group comprising a solubilizing agent, an absorption promoter, a film-forming agent, a gelling agent or their mixtures,

10 combined with or mixed with suitable excipients for the production of a gelled and/or film-forming pharmaceutical form.

2. Topical hormonal composition with systemic effect according to claim 1, characterized in that the progestogen derived from 19-nor progesterone is nomegestrol and/or one of its esters or
15 ethers.

3. Topical hormonal composition with systemic effect according to claim 1 or claim 2, characterized in that the progestogen derived from 19-nor progesterone is nomegestrol acetate.

20 4. Topical hormonal composition with systemic effect according to any one of the claims 1 to 3, characterized in that the quantity of nomegestrol or of one of its esters or ethers varies from 0.05 to 1% by weight of the total composition.

25 5. Topical hormonal composition with systemic effect according to claim 4, characterized in that the quantity of nomegestrol or of one of its esters or ethers is 0.4 % by weight of the total composition.

30 6. Topical hormonal composition with systemic effect according to any one of the claims 1 to 5, characterized in that the solubilizing agent is chosen from the group comprising water, alcohols, propyleneglycol, a C₈/C₁₀ polyoxyethylene glycosyl glyceride or their mixtures.

7. Topical hormonal composition with systemic effect according to any one of the claims 1 to 6, characterized in that the solubilizing agent is a ternary mixture of 95° ethanol / water /

propyleneglycol, in which the percentage of 95° ethanol varies from 30 to 50 %, that of water from 30 to 60 % and that of propyleneglycol from 2 to 20 % .

8. Topical hormonal composition with systemic effect according to any one of the claims 1 to 6, characterized in that the solubilizing agent is a quaternary mixture 95° ethanol / water / Labrasol® / propyleneglycol, in which the percentage of 95° ethanol varies from 30 to 50 %, that of water from 30 to 60 %, that of Labrasol® from 3 to 7 % and that of propyleneglycol from 2 to 20 %.

9. Topical hormonal composition with systemic effect according to any one of the claims 1 to 8, characterized in that the absorption promoter is chosen from the group consisting of isopropylideneglycerol, α -tocopheryl polyethyleneglycol 1000 succinate and monoethyl ether of diethylene glycol.

10. Topical hormonal composition with systemic effect according to claim 9, characterized in that the absorption promoter is isopropylideneglycerol.

11. Topical hormonal composition with systemic effect according to any one of the claims 1 to 10, characterized in that the gelling agent is chosen from the group consisting of cellulose derivatives and acrylic derivatives.

12. Topical hormonal composition with systemic effect according to claim 11, characterized in that the cellulose derivative is hydroxypropylmethylcellulose.

13. Topical hormonal composition with systemic effect according to claim 11, characterized in that the acrylic derivative is a carbomer.

14. Topical hormonal composition with systemic effect according to any one of the claims 1 to 13, characterized in that the film-forming agent is chosen from the group consisting of cellulose derivatives, methacrylic derivatives and polyvinylpyrrolidone derivatives.

15. Topical hormonal composition with systemic effect according to claim 14, characterized in that the cellulose derivative is hydroxypropylmethylcellulose acetate succinate.

16. Topical hormonal composition with systemic effect according to claim 14, characterized in that the methacrylic derivative is an aqueous dispersion of an anionic copolymer of methacrylic acid and ethyl acrylate.

5 17. Topical hormonal composition according to any one of the claims 1 to 16, characterized in that it is in the form of a gel or a film-forming gel and in that it contains, in an aqueous-alcoholic mixture, 8 % of propyleneglycol and 3 % of isopropylidene glycerol.

ABSTRACT

The present invention relates to the area of chemotherapy and more especially to the development of new galenic forms for application to the skin.

5

It relates more particularly to a topical hormonal composition with systemic action for the correction of progesterone deficiency in premenopausal women and for hormone replacement in menopausal women, characterized in that it contains, as the active principle, a progestogen derived from 19-nor progesterone, a vehicle permitting systemic passage of the said active principle chosen from the group comprising a solubilizing agent, an absorption promoter, a film-forming agent, a gelling agent or their mixtures, combined with or mixed with suitable excipients for production of a pharmaceutical form as a gel and/or a film.

10

09/646763

1/6

Figure 1





Figure 2



09/646763

2/6

Figure 3





Figure 4



09/646763

3/6

Figure 5





Figure 6



09/646763

4/6

Figure 7





Figure 8



09/646763

5/6

Figure 9





Figure 10



09/646763

6/6

Figure 11





Figure 12



Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (8-56)
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
For use through 9/30/98. OMB 0651-0032
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION

ADDITIONAL INVENTOR(S) Supplemental Sheet

Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	JACQUES	Middle Initial		Family Name	PARIS	Suffix	
Inventor's Signature	JACQUES PARIS				Date	10.03.2000	
Residence: City	Nice	State	FRX	Country	France	Citizenship	FR
Post Office Address							
Post Office Address							
City	Nice	State		Zip	F-06100	Country	France
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	JEAN-LOUIS	Middle Initial		Family Name	THOMAS	Suffix	
Inventor's Signature	JEAN LOUIS THOMAS				Date	10.03.2000	
Residence: City	Charenton Le Pont	State	FRX	Country	France	Citizenship	FR
Post Office Address							
Post Office Address							
City	Charenton Le Pont	State		Zip	F-94220	Country	France
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name		Middle Initial		Family Name		Suffix	
Inventor's Signature					Date		
Residence: City		State		Country		Citizenship	
Post Office Address							
Post Office Address							
City		State		Zip		Country	
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name		Middle Initial		Family Name		Suffix	
Inventor's Signature					Date		
Residence: City		State		Country		Citizenship	
Post Office Address							
Post Office Address							
City		State		Zip		Country	
<input type="checkbox"/> Additional inventors are being named on supplemental sheet(s) attached hereto							

GEI-078

Please print a title page for this form.

[4]

Approved for use through 9/30/99. OMB 051-0022
 Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE
 Under the Patent and Trademark Act of 1980, no person is required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or §365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Name	Registration Number	Name	Registration Number
Charles A. Muserlian	19,683		
Jordan B. Bierman	18,629		
Donald C. Lucas	31,275		
Bierman, Muserlian and Lucas	18,818		

☐ Additional registered practitioner(s) named on a supplemental sheet attached hereto.

Direct all correspondence to:

Name	Bierman, Muserlian and Lucas		
Address			
Address	600 Third Avenue		
City	New York	State	New York
Country	U.S.A.	Telephone	(212) 661-8000
		Fax	(212) 661-8002

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like, so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:

☐ A petition has been filed for this unsigned inventor

Given Name	MICHEL	Middle Initial		Family Name	LANQUETIN	Suffix e.g. Jr.	
Inventor's Signature	MICHEL LANQUETIN				Date	10/03/2000	

Residence City	La Trinite	State	FR	Country	France	Citizenship	FR
Post Office Address							
Post Office Address	Chemin Soanes, Quartier de l'Adrech						
City	La Trinite	State		Zip	P-06340	Country	France

☒ Additional inventors are being named on supplemental sheet(s) attached hereto

Please type a plus sign (+) inside this box

PTO/SB/01 (8-96)
 Approved for use through 9/30/98. OMB 0651-0032
 Patent and Trademark Office U.S. DEPARTMENT OF COMMERCE
 Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION

☒ Declaration Submitted with Initial Filing OR ☐ Declaration Submitted after Initial Filing

Attorney Docket Number	GEI-078
First Named Inventor	LANGUETIN et al
COMPLETE IF KNOWN	
Application Number	PCT/FR99/00680
Filing Date	3/23/99
Group Art Unit	
Examiner Name	

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

TOPICAL HORMONAL COMPOSITION WITH SYSTEMIC ACTION

(Title of the invention)

the specification of which

☐ is attached hereto OR

☒ was filed on (MM/DD/YYYY) 3/23/99 as United States Application Number or PCT International

Application Number PCT/FR99/00680 and was amended on (MM/DD/YYYY) (if applicable)

I hereby state that I have reviewed and understand the contents of the above identifying specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35 United States Code § 119 (a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365 (a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
98/03533	France	3/23/98	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
PCT/FR99/00680	France	3/23/99	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority sheet attached hereto.

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority sheet attached hereto.

(Page 1 of 5)

Under Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner of Patents and Trademarks, Washington, DC 20231

(January 1997)